

Original Article

Hepatic steatosis causes induction of the chemokine RANTES in the absence of significant hepatic inflammation

Georgi Kirovski^{1,*}, Erwin Gäbele^{1,*}, Christoph Dorn¹, Lukas Moleda¹, Christoph Niessen¹, Thomas S. Weiss², Hella Wobser¹, Doris Schacherer¹, Christa Buechler¹, Hermann E. Wasmuth³, Claus Hellerbrand¹

¹Department of Internal Medicine I, University Hospital Regensburg, Regensburg, Germany; ²Center for Liver Cell Research, Department of Surgery, University Hospital Regensburg, Regensburg, Germany; ³Department of Medicine III, University Hospital Aachen, Aachen, Germany. *Equal contributors.

Received July 16, 2010; accepted July 31, 2010; available online August 2, 2010

Abstract: Nonalcoholic fatty liver disease (NAFLD) encompasses a spectrum ranging from simple steatosis to cirrhosis. Hepatocellular lipid accumulation is a hallmark of both nonalcoholic steatosis and steatohepatitis (NASH). The latter develops upon pro-inflammatory cell infiltration and is widely considered as the first relevant pathophysiological step in NAFLD-progression. The chemokine CCL5/RANTES plays an important role in the progression of hepatic inflammation and fibrosis. We here aimed to investigate its expression in NAFLD. Incubation of primary human hepatocytes with palmitic acid induced a dose-dependent lipid accumulation, and corresponding dose-dependent RANTES induction *in vitro*. Furthermore, we observed significantly elevated hepatic RANTES expression in a dietary model of NAFLD, in which mice were fed a high-fat diet for 12 weeks. This diet induced significant hepatic steatosis but only minimal inflammation. In contrast to the liver, RANTES expression was not induced in visceral adipose tissue of the group fed with high-fat diet. Finally, RANTES serum levels were elevated in patients with ultrasound-diagnosed NAFLD. In conclusion, our data indicate hepatocytes as cellular source of elevated hepatic as well as circulating RANTES levels in response to hepatic steatosis. Noteworthy, upregulation of RANTES in response to lipid accumulation occurs in the absence of relevant inflammation, which further indicates that hepatic steatosis *per se* has pathophysiological relevance and should not be considered as benign.

Keywords: Hepatic steatosis, liver cirrhosis, nonalcoholic steatosis, steatohepatitis, chemokine CCL5, RANTES, inflammation, nonalcoholic fatty liver disease (NAFLD)

Introduction

Non-alcoholic fatty liver disease (NAFLD) is a condition with excessive fat accumulation in hepatocytes [1]. NAFLD is often classified as primary and secondary according to the underlying etiology [2]. Primary NAFLD generally refers to fatty liver associated with features of the metabolic syndrome [3], and in fact, NAFLD may be considered as hepatic manifestation of the metabolic syndrome [4].

Recent studies on the development and progression of NAFLD emphasize the pathophysiological role of cytokines and chemokines besides that of insulin resistance, oxidative stress and lipid peroxidation [5]. Chemokines repre-

sent a large superfamily of proteins that possess diverse biologic activities [6]. Regulated upon Activation, Normal T-cell Expressed, and Secreted (RANTES, also known as CCL5) is a CC beta-chemokine with strong chemoattractant activity for T lymphocytes and monocytes [7]. It has been implicated in the pathogenesis of numerous diseases including cancer [8], neurological disorders [9], asthma [10], autoimmune diseases such as rheumatoid arthritis [11] and renal diseases [12].

Increased (hepatic) RANTES expression has been reported in patients with chronic viral hepatitis [13,14], and a comprehensive study by Seki and colleagues [15] has shown that (increased expression of) RANTES promotes

hepatic inflammation and fibrosis in the bile duct ligation model and in a model of chronic toxic liver injury by application of tetrachloro-carbon. Furthermore, it has been reported that obesity promotes chemokine expression in patients with chronic hepatitis C infection [16], and increased hepatic RANTES expression has been observed in a murine NASH-model [17] and in obese patients [18].

Based on the hypothesis that increased hepatic RANTES expression is an early event in NAFLD and a critical factor influencing its progression the aim of the present study was to analyze the effect of hepatocellular lipid accumulation on RANTES expression in the absence of significant hepatic inflammation.

Materials and methods

Primary human hepatocytes and in vitro model of hepatocellular lipid accumulation

Isolation and culture of primary human hepatocytes (PHH) were performed as described [19]. Human liver tissue for cell isolation was obtained according to the guidelines of the charitable state-controlled foundation HTCR with the patient's informed consent. Hepatocellular lipid accumulation was induced as described [20]. Briefly, a 10 mM palmitate/1% bovine serum albumin (BSA) stock solution was obtained by complexing the appropriate amount of palmitate to BSA. PHH were grown in DMEM supplemented with penicillin (400 U/ml), streptomycin (50 mg/ml), L-glutamine (300 mg/ml), 0.2% FCS and palmitate for 24h in different concentrations as indicated. FFA-free-BSA-treated cells served as controls.

Intracellular triglyceride assay

Total triglycerides were extracted using the method of Bligh and Dyer with slight modifications [21,22] and quantified by the triglyceride determination kit (GPO) (Sigma, Deisenhofen, Germany) as described [20].

Murine model of hepatic steatosis

Male BALB/c mice were purchased from Charles River Laboratories (Sulzfeld, Germany) at 6 weeks of age and housed under standard conditions. After 1 week of acclimatization mice were divided into two groups (n=6 each) and

fed either with control diet or a high fat diet (HFD) containing 30% lard. All chows were prepared by Ssniff (Soest, Germany). After 12 weeks feeding animals were sacrificed. Liver tissue and epididymal fat were immediately snap-frozen and stored at -80 °C for subsequent analysis.

Human serum samples of patients with ultrasound-diagnosed fatty liver and control patients

Serum samples were obtained from randomly selected patients recruited from the referrals of the ultrasound (US) centre at our clinic from January 2008 to January 2009. According to the results of a standardized ultrasound examination participants were divided into 2 groups – a control group with normal US liver appearance and a group of patients with US-diagnosed fatty liver. The following exclusion criteria were applied: history of malignancy of any kind, chronic hepatobiliary diseases, use of medication known to affect hepatic steatosis (e.g. glucocorticoids, anti-estrogens), inflammatory bowel disease, human immunodeficiency virus infection, drug or alcohol abuse (more than 20 g/day), familial hyperlipidemia, and acute medical emergencies. Further, patients with clinical or laboratory signs of systemic inflammation were excluded to omit possible confounding effect on serum RANTES.

After collection serum samples were immediately snap frozen and stored at -80 °C. Informed consent was obtained from all patients and the study was approved by the local Ethics Committee.

RANTES expression analysis

Isolation of total cellular RNA from cultured cells and tissues and reverse transcription were performed as described [23]. Quantitative real time-PCR was performed with specific sets of primers applying LightCycler technology (Roche, Mannheim, Germany) as described [24] and specific human and mouse RANTES Primer assays (Qiagen, Hilden, Germany).

Analysis of RANTES serum levels

RANTES in human sera was assessed by sandwich enzyme-linked immunosorbent assay (ELISA) using the human CCL5 DuoSet ELISA development kit (R&D Systems Inc., MN, USA, catalog number DY278).

RANTES in hepatic steatosis




Figure 1. RANTES expression in an *in vitro* model of hepatocellular lipid accumulation. Primary human hepatocytes were incubated with 0.4 μM, 0.6 μM and 0.8 μM palmitate (Palm.) for 24 h, which leads to a dose dependent intracellular lipid accumulation as shown before [20]. Subsequently, RNA was isolated, reverse transcribed and RANTES mRNA expression was analyzed by quantitative PCR. (* $p < 0.05$ compared to control (ctr.)).

Statistical analysis

Statistical analysis was performed using SPSS version 15.0 (SPSS, Chicago, IL, USA) and GraphPad Prism Software (GraphPad Software, Inc., San Diego, USA). Results are expressed as means \pm standard error (range). Comparisons

between groups were made using one way analysis of variance (with Bonferroni correction for multiple comparisons). P-values < 0.05 were considered as statistically significant.

Results

Hepatocellular lipid accumulation induced RANTES expression *in vitro*

Recently, we have described an *in vitro* model of hepatocellular lipid accumulation [20]. Incubation of primary human hepatocytes (PHH) with palmitic acid led to a dose dependent increase of hepatocellular free fatty acid and triglyceride levels [20]. Interestingly, this lipid accumulation led to a significant induction of RANTES mRNA expression in PHH (**Figure 1**).

Increased hepatic RANTES expression in steatotic murine livers

Next, we analyzed RANTES expression in livers of mice fed a high-fat diet (HFD) [25]. Feeding this diet led to significant hepatic lipid accumulation (**Figure 2A**). In contrast mice in the HFD-group had normal levels serum transaminases (**Figure 2B**) and revealed only slightly elevated TNF mRNA expression (**Figure 2C**).

Activated hepatic stellate cells (HSC) are the cellular source of excessive extracellular matrix




Figure 2. RANTES expression in the liver and in visceral fat in a murine model of NAFLD. Mice were fed a high fat diet leading to hepatic steatosis. Mice receiving standard chow served as control (ctr.). **(A)** Assessment of hepatic triglyceride content. **(B)** Serum GOT levels. Analysis of **(C)** hepatic TNF, collagen I, α -smooth muscle actin (α -sma), and RANTES mRNA expression, and **(D)** expression of RANTES mRNA in visceral fat by quantitative PCR. (* $p < 0.05$ compared to control).

deposition in chronic liver injury [26], and previous studies have shown increased RANTES expression in activated HSC [27]. However, neither expression of alpha-smooth muscle actin, a sensitive marker of HSC activation, nor collagen I expression was elevated in HFD-fed mice, indicating that there was no relevant activation of HSC in the livers of these animals (Figure 2C). However and notably, RANTES expression was significantly increased in mice fed with the HFD (Figure 2C).

It has been shown that visceral fat is a significant source of proinflammatory cytokines including RANTES [28,29]. Therefore, we also analyzed RANTES expression in visceral fat but found only minimally increased mRNA expression compared to control mice (Figure 2D).

Increased RANTES serum levels in patients with non-alcoholic fatty liver

Recently, we have described a cohort of patients with referral for sonographic examination of the abdomen and ultrasound-diagnosed NAFLD [30]. Analysis of serum levels of RANTES revealed slightly but significantly elevated RANTES levels in patients with ultrasound-diagnosed NAFLD (n=45; mean: 85.9 ng/ml, SEM 3.7 ng/ml,) in comparison to the control group (n=61; mean: 60.1 ng/ml, SEM 2.9 ng/ml; p<0.001), (Figure 3).

Discussion

The chemokine RANTES has been shown to play a central role in the pathogenesis and progression of chronic liver disease, however, present information regarding its expression in nonalcoholic fatty liver disease (NAFLD) is sparse [31].

RANTES has been shown to be expressed by hepatocytes and is up-regulated in response to inflammatory conditions [14;32-34]. Here, we reveal that lipid accumulation did also dose-dependently induce RANTES expression in primary human hepatocytes *in vitro*. Thus, pure hepatocellular steatosis leads to increased hepatic RANTES expression. In line with this, we observed that feeding a high-fat diet induced hepatic RANTES expression in the absence of significant hepatic inflammation or fibrosis. This finding further confirms steatotic hepatocytes but not infiltrating inflammatory cells or activated hepatic stellate cells as main cellular sources of elevated hepatic RANTES expression




Figure 3. Serum RANTES levels in patients with NAFLD. Serum RANTES levels in patients with ultrasound (US) diagnosed fatty liver compared to a control (ctr.) group of patients with normal sonographic liver appearance. (*p<0.05 compared to control).

in high-fat diet fed mice.

Nevertheless, it has to be considered that obesity frequently leads to up-regulation of cytokines as a part of a systemic state of low inflammation not only in the liver but also in (visceral) adipose tissue, which is another important source of circulating cytokines including RANTES [29]. However, in our study, RANTES expression in visceral fat of mice, which were fed with a high-fat diet, was similar to the control group. This suggests that at least under these experimental conditions the fatty liver but not adipose tissue is the source of elevated (circulating) RANTES levels.

Taken together with our *in vitro* data and the known pathophysiological role of RANTES in other chronic liver diseases, this indicates that steatosis-triggered RANTES production by hepatocytes is an early event during the natural course of NAFLD and may be involved, along with other factors/cytokines, in the progression of fatty liver to significant inflammation, e.g. NASH.

Interestingly, patients with US-diagnosed NAFLD had significantly elevated RANTES serum levels

compared to the control group, although this increase was only moderate compared to the induction of RANTES mRNA expression in *in vitro* lipid loaded hepatocytes and murine fatty livers. However, although US examination can be adequately used to assess hepatic steatosis [35,36], it has to be considered that the reliable threshold is approximately 30% fat in the hepatic tissue [37]. Thus, it is likely that a significant number of patients in the control group also had hepatic steatosis below the US-detection level. NAFLD is generally defined by lipid deposition greater than 5-10% of the liver weight [1], and our *in vitro* data indicate that even minimal hepatocellular lipid accumulation leads to increased RANTES levels. Conversely, despite the lack of histological examination in this hospital cohort of randomly selected patients, it can be estimated from epidemiological studies [38], that only a minority of cases had significant hepatic inflammation (e.g. criteria for NASH), which may have caused elevated RANTES levels.

In summary, our study indicates that hepatic steatosis causes an up-regulation of hepatic RANTES expression. Noteworthy, increased expression of RANTES in response to hepatocellular lipid accumulation can occur in the absence of relevant hepatic inflammation. This finding further indicates that hepatic steatosis *per se* has pathophysiological relevance and should not be considered as benign.

Acknowledgement

We want to thank Marina Fink and Birgitta Ott-Rötzer for excellent technical assistance, and Hanna Huber and Catrin Beer for collecting sera of patients.

This work was supported by grants from the German Research Association (DFG) to C.H. and the Medical Faculty of the University of Regensburg (ReForM) to E.G., T.S.W., C.B. and C.H. H.E.W is supported by the DFG (SFB-TRR 57).

Please address all correspondences to: Claus Hellerbrand, MD, University of Regensburg, Department of Internal Medicine I, D-93042 Regensburg, Germany, Tel: +49-941-944-7155, Fax: +49-941-944-7154, E-mail: claus.hellerbrand@klinik.uni-regensburg.de

References

[1] Neuschwander-Tetri BA and Caldwell SH. Non-

alcoholic steatohepatitis: summary of an AASLD Single Topic Conference. *Hepatology* 2003; 37: 1202-1219.

[2] Angulo P. Nonalcoholic fatty liver disease. *N Engl J Med* 2002; 346: 1221-1231.

[3] Clark JM. The epidemiology of nonalcoholic fatty liver disease in adults. *J Clin Gastroenterol* 2006; 40 Suppl 1: S5-10.

[4] Marchesini G, Brizi M, Bianchi G, Tomassetti S, Bugianesi E, Lenzi M, McCullough AJ, Natale S, Forlani G, and Melchionda N. Nonalcoholic fatty liver disease: a feature of the metabolic syndrome. *Diabetes* 2001; 50: 1844-1850.

[5] Jiang J and Torok N. Nonalcoholic steatohepatitis and the metabolic syndrome. *Metab Syndr Relat Disord* 2008; 6: 1-7.

[6] Coelho AL, Hogaboam CM, and Kunkel SL. Chemokines provide the sustained inflammatory bridge between innate and acquired immunity. *Cytokine Growth Factor Rev* 2005; 16: 553-560.

[7] Appay V and Rowland-Jones SL. RANTES: a versatile and controversial chemokine. *Trends Immunol* 2001; 22: 83-87.

[8] Soria G and Ben-Baruch A. The inflammatory chemokines CCL2 and CCL5 in breast cancer. *Cancer Lett* 2008; 267: 271-285.

[9] Szczucinski A and Losy J. Chemokines and chemokine receptors in multiple sclerosis. Potential targets for new therapies. *Acta Neurol Scand* 2007; 115: 137-146.

[10] Bisset LR and Schmid-Grendelmeier P. Chemokines and their receptors in the pathogenesis of allergic asthma: progress and perspective. *Curr Opin Pulm Med* 2005; 11: 35-42.

[11] Yang MH, Wu FX, Xie CM, Qing YF, Wang GR, Guo XL, Tang Z, Zhou JG, and Yuan GH. Expression of CC chemokine ligand 5 in patients with rheumatoid arthritis and its correlation with disease activity and medication. *Chin Med Sci J* 2009; 24: 50-54.

[12] Krensky AM and Ahn YT. Mechanisms of disease: regulation of RANTES (CCL5) in renal disease. *Nat Clin Pract Nephrol* 2007; 3: 164-170.

[13] Nischalke HD, Nattermann J, Fischer HP, Sauerbruch T, Spengler U, and Dumoulin FL. Semi-quantitative analysis of intrahepatic CC-chemokine mRNAs in chronic hepatitis C. *Mediators Inflamm* 2004; 13: 357-359.

[14] Zeremski M, Petrovic LM, and Talal AH. The role of chemokines as inflammatory mediators in chronic hepatitis C virus infection. *J Viral Hepat* 2007; 14: 675-687.

[15] Seki E, de MS, Gwak GY, Kluwe J, Inokuchi S, Bursill CA, Llovet JM, Brenner DA, and Schwabe RF. CCR1 and CCR5 promote hepatic fibrosis in mice. *J Clin Invest* 2009; 119: 1858-1870.

[16] Palmer C, Corpuz T, Guirguis M, O'Toole S, Yan K, Bu Y, Jorgenson J, Talbot M, Loi K, Lloyd A, and Zekry A. The effect of obesity on intrahepatic cytokine and chemokine expression in

- chronic hepatitis C infection. *Gut* 2010; 59: 397-404.
- [17] Desai MS, Mariscalco MM, Tawil A, Vallejo JG, and Smith CW. Atherogenic diet-induced hepatitis is partially dependent on murine TLR4. *J Leukoc Biol* 2008; 83: 1336-1344.
- [18] Wu H, Ghosh S, Perrard XD, Feng L, Garcia GE, Perrard JL, Sweeney JF, Peterson LE, Chan L, Smith CW, and Ballantyne CM. T-cell accumulation and regulated on activation, normal T cell expressed and secreted upregulation in adipose tissue in obesity. *Circulation* 2007; 115: 1029-1038.
- [19] Weiss TS, Jahn B, Cetto M, Jauch KW, and Thasler WE. Collagen sandwich culture affects intracellular polyamine levels of human hepatocytes. *Cell Prolif* 2002; 35: 257-267.
- [20] Wobser H, Dorn C, Weiss TS, Amann T, Bollheimer C, Buttner R, Scholmerich J, and Hellerbrand C. Lipid accumulation in hepatocytes induces fibrogenic activation of hepatic stellate cells. *Cell Res* 2009; 19: 996-1005.
- [21] Novo E, Marra F, Zamara E, Valfre di BL, Monitillo L, Cannito S, Petrai I, Mazzocca A, Bonacchi A, De Franco RS, Colombatto S, Autelli R, Pinzani M, and Parola M. Overexpression of Bcl-2 by activated human hepatic stellate cells: resistance to apoptosis as a mechanism of progressive hepatic fibrogenesis in humans. *Gut* 2006; 55: 1174-1182.
- [22] Schnabl B, Choi YH, Olsen JC, Hagedorn CH, and Brenner DA. Immortal activated human hepatic stellate cells generated by ectopic telomerase expression. *Lab Invest* 2002; 82: 323-333.
- [23] Hellerbrand C, Mühlbauer M, Wallner S, Schuierer M, Behrmann I, Bataille F, Weiss T, Schölmerich J, and Bosserhoff AK. Promoter-hypermethylation is causing functional relevant downregulation of methylthioadenosine phosphorylase (MTAP) expression in hepatocellular carcinoma. *Carcinogenesis* 2006; 27: 64-72.
- [24] Mühlbauer M, Bosserhoff AK, Hartmann A, Thasler WE, Weiss TS, Herfarth H, Lock G, Scholmerich J, and Hellerbrand C. A novel MCP-1 gene polymorphism is associated with hepatic MCP-1 expression and severity of HCV-related liver disease. *Gastroenterology* 2003; 125: 1085-1093.
- [25] Dorn C, Riener MO, Kirovski G, Saugspier M, Steib K, Weiss TS, Gäbele E, Kristiansen G, Hartmann A, and Hellerbrand C. Expression of fatty acid synthase in nonalcoholic fatty liver disease. *Int J Clin Exp Pathol* 2010; 3: 505-514.
- [26] Parola M, Marra F, and Pinzani M. Myofibroblast-like cells and liver fibrogenesis: Emerging concepts in a rapidly moving scenario. *Mol Aspects Med* 2008; 29: 58-66.
- [27] Schwabe RF, Batailler R, and Brenner DA. Human hepatic stellate cells express CCR5 and RANTES to induce proliferation and migration. *Am J Physiol Gastrointest Liver Physiol* 2003; 285: G949-G958.
- [28] Madani R, Karastergiou K, Ogston NC, Miheisi N, Bhome R, Haloob N, Tan GD, Karpe F, Malone-Lee J, Hashemi M, Jahangiri M, and Mohamed-Ali V. RANTES release by human adipose tissue in vivo and evidence for depot-specific differences. *Am J Physiol Endocrinol Metab* 2009; 296: E1262-E1268.
- [29] Skurk T, Mack I, Kempf K, Kolb H, Hauner H, and Herder C. Expression and secretion of RANTES (CCL5) in human adipocytes in response to immunological stimuli and hypoxia. *Horm Metab Res* 2009; 41: 183-189.
- [30] Kirovski G, Schacherer D, Wobser H, Huber H, Niessen C, Beer C, Schoelmerich J and Hellerbrand C. Prevalence of ultrasound-diagnosed non-alcoholic fatty liver disease in a hospital cohort and its association with anthropometric, biochemical and sonographic characteristics. *Int J Clin Exp Med* 2010 [in press]
- [31] Wasmuth HE, Tacke F and Trautwein C. Chemokines in liver inflammation and fibrosis. *Sem Liv Dis* 2010 [in press]
- [32] Afford SC, Fisher NC, Neil DA, Fear J, Brun P, Hubscher SG, and Adams DH. Distinct patterns of chemokine expression are associated with leukocyte recruitment in alcoholic hepatitis and alcoholic cirrhosis. *J Pathol* 1998; 186: 82-89.
- [33] Rowell DL, Eckmann L, Dwinell MB, Carpenter SP, Raucy JL, Yang SK, and Kagnoff MF. Human hepatocytes express an array of proinflammatory cytokines after agonist stimulation or bacterial invasion. *Am J Physiol* 1997; 273: G322-G332.
- [34] Ohashi T, Tanabe J, Ishikawa T, Okumura A, Sato K, Ayada M, Hotta N, Kuzuya T, Ito H, Nakao H, Yoneda M, and Kakumu S. Inflammatory cytokines modulate chemokine production patterns of HepG2 cells toward initially inclined direction. *Hepatol Res* 2009; 39: 510-519.
- [35] Joseph AE, Saverymuttu SH, al-Sam S, Cook MG, and Maxwell JD. Comparison of liver histology with ultrasonography in assessing diffuse parenchymal liver disease. *Clin Radiol* 1991; 43: 26-31.
- [36] Saverymuttu SH, Joseph AE, and Maxwell JD. Ultrasound scanning in the detection of hepatic fibrosis and steatosis. *Br Med J (Clin Res Ed)* 1986; 292: 13-15.
- [37] Saadeh S, Younossi ZM, Remer EM, Gramlich T, Ong JP, Hurley M, Mullen KD, Cooper JN, and Sheridan MJ. The utility of radiological imaging in nonalcoholic fatty liver disease. *Gastroenterology* 2002; 123: 745-750.
- [38] Bellentani S, Scaglioni F, Marino M, and Bedogni G. Epidemiology of non-alcoholic fatty liver disease. *Dig Dis* 2010; 28: 155-161.