

## Original Article

# ***Propionibacterium acnes* induces intervertebral discs degeneration by increasing MMP-1 and inhibiting TIMP-1 expression via the NF- $\kappa$ B pathway**

Yuehuan Zheng<sup>1\*</sup>, Yazhou Lin<sup>2,3\*</sup>, Zhe Chen<sup>2,3</sup>, Yucheng Jiao<sup>2,3</sup>, Ye Yuan<sup>2,3</sup>, Changwei Li<sup>3</sup>, Xiangyang Xu<sup>2</sup>, Peng Cao<sup>2,3</sup>

<sup>1</sup>Department of Orthopedics, Ruijin Hospital North, Shanghai Jiaotong University School of Medicine, Shanghai, China; <sup>2</sup>Department of Orthopedics, Ruijin Hospital, Shanghai Jiaotong University School of Medicine, Shanghai, China; <sup>3</sup>Shanghai Key Laboratory for Prevention and Treatment of Bone and Joint Diseases with Integrated Chinese-Western Medicine, Shanghai Institute of Traumatology and Orthopedics, Ruijin Hospital, Shanghai Jiaotong University School of Medicine, Shanghai, China. \*Equal contributors.

Received April 22, 2018; Accepted May 23, 2018; Epub July 1, 2018; Published July 15, 2018

**Abstract:** Emerging evidence suggests that *Propionibacterium acnes* (*P. acnes*) may be a new pathogen implicated in intervertebral disc degeneration (IVDD), although the underlying mechanisms are unclear. Since the most significant biochemical change of IVDD is inability of the extracellular matrix (ECM) synthesis by nucleus pulposus cells, we first analyzed the expression of aggrecan and collagen II in nucleus pulposus tissues of IVDD patients with or without *P. acnes* infection. Compared with the *P. acnes*-negative controls, the expression levels of aggrecan and collagen II were significantly dampened in nucleus pulposus tissues with *P. acnes* infection. Interestingly, we found that *P. acnes* infection strongly increased matrix-degrading metalloproteinase-1 (MMP-1) expression but decreased that of tissue inhibitor of metalloproteinase-1 (TIMP1). Furthermore, the dampened aggrecan and collagen II synthesis concomitant with the increased MMP-1 and decreased TIMP-1 expression seen in *P. acnes*-induced IVDD were confirmed by a rat model. Mechanistically, *P. acnes* infection increased MMP-1 levels, while decreasing TIMP-1 expression via the NF- $\kappa$ B pathway. Overall, these findings reveal that *P. acnes* infection dampens aggrecan and collagen II synthesis in nucleus pulposus cells by increasing MMP-1 and inhibiting TIMP-1 expression via the NF- $\kappa$ B pathway, which may ultimately lead to IVDD.

**Keywords:** *Propionibacterium acnes*, MMP-1, TIMP-1, NF- $\kappa$ B, Disc degeneration

## Introduction

Intervertebral discs (IVDs) are complex anatomical structures with a specific load-bearing organization that gives the spine flexibility [1]. As the inner core of IVDs, the nucleus pulposus (NP) is known to resist mechanical loading by synthesizing the extracellular matrix (ECM) containing proteoglycan and collagen II, thus maintaining the stability of IVDs [2]. The hallmark of progressive IVD degeneration (IVDD) is the incapacity of NP cells (NPCs) to maintain normal homeostatic tissue remodeling, thereby leading to a series of severe sequelae, such as disc herniation, Modic changes, sciatica, and lower back pain, which greatly worsen quality of life and increase the economic and health care burdens of society [3].

Traditionally, excessive mechanical loading, nutritional disorders, traumatic injury, or genetic predisposition are considered the main etiological causes for IVDD [3]. However, recent studies suggested that *Propionibacterium acnes*, a low-virulence anaerobic bacterium, which was found to latently reside inside non-pyogenic IVDs, could induce IVDD. Epidemiological studies and systemic analyses revealed that the prevalence of *P. acnes* in IVDD ranged from 13 to 44% [4-8]. In our previous study, *P. acnes* colonies were identified in non-pyogenic degenerated IVD by anaerobic culture and histological observation [9]. More interestingly, inoculation of *P. acnes* into normal rabbit IVDs induced severe disc degeneration and Modic changes [10]. Although our previous study uncovered that *P. acnes* infection could initiate IVDD by

## *P. acnes* induces disc degeneration via MMP-1 and TIMP-1

promoting NPC apoptosis [11], the pathological mechanisms of *P. acnes*-induced IVDD remain unclear and need further investigation.

In the present study, we set out to investigate the mechanisms by which *P. acnes* induces IVDD. Our findings uncover that *P. acnes* infection dampens aggrecan and collagen II synthesis in nucleus pulposus cells by increasing MMP-1 and inhibiting TIMP-1 expression via the NF- $\kappa$ B pathway, which may ultimately lead to IVDD.

### Materials and methods

#### *Patients and tissue harvesting*

A total of 20 patients were included in this study. The patients underwent discectomy at the single-level lumbar spine due to disc degeneration associated with low back pain and/or sciatica. Patients who received antibiotics within the month preceding surgery were not included in this study. Following the methodology of previous research [9], all tissues were cultured in tryptone soy broth (TSB) for 14 days under anaerobic conditions, then the presence of bacteria in the culture was identified by amplifying the 16S rDNA gene by PCR according to our previous protocol [9].

After culture of the specimen and bacterial identification, the patients who had *P. acnes* only in intervertebral discs were classified as the positive group (n=23). A case-controlled method was used for the quantitative analysis following a previous study [12]. 10 patients who were identified as completely bacteria-free in their intervertebral discs were selected to match each of the positive patients based on the following criteria: 1. same gender; 2. same surgery segment; 3. same symptoms of low back pain only, sciatica only or both; 4. similar ages  $\pm$  5 years; 5. similar duration of symptoms  $\pm$  3 months.

The study was approved by the Institutional Review Board of Shanghai Ruijin Hospital and informed consent forms were signed by all patients.

#### *Preparation of P. acnes inoculum*

*P. acnes* preparation was performed as described in our previous report [11]. A standard strain of *P. acnes* (ATCC: 6919, GIM:

1.243, Guangdong Microbiology Culture Center, Guangdong, China) was cultured on Gifu Anaerobic (GAM) broth (Nissui, Tokyo, Japan) for 3 d at 37°C under anaerobic conditions.

#### *Inoculation of P. acnes into rat caudal intervertebral discs*

The inoculation of *P. acnes* into caudal rat IVDs was performed as described in our previous study [11]. The target vertebrae (Ca) 6/7 to (Ca) 8/9 (n=3 per animal) of eight-week-old male Sprague-Dawley rats were identified and marked by palpation and X-ray before surgery. A volume of 2.5  $\mu$ l *P. acnes* (OD<sub>600</sub>=3.0), *P. acnes* with BAY11-7082 (NF- $\kappa$ B inhibitor, 10  $\mu$ M) or saline was inoculated vertically into the nucleus pulposus using a microsyringe with a 28-gauge needle (Hamilton, Nevada, USA). All animal experiments were performed in accordance with the protocol approved by the Shanghai Jiao Tong University (SJTU) Animal Care and Use Committee [IACUC protocol number: SYXK (Shanghai) 2011-0113] and in accordance with the Ministry of Science and Technology of the People's Republic of China animal care guidelines. All surgeries were performed under anesthesia, and all efforts were made to minimize suffering.


#### *Cell culture and stimulation*

Primary nucleus pulposus cells were isolated and cultured as described in our previous report [11]. The bacteria were harvested from 3-day cultures in a stationary phase and washed twice with phosphate-buffered saline (PBS). The bacterial density was adjusted to optical density (OD<sub>600</sub>=2). Then, *P. acnes* were added to the cell culture ( $5 \times 10^5$  cells/well) in a 6-well culture plate at a 100:1 multiplicity of infection (MOI) without antibiotics.

#### *Immunoblotting and immunofluorescent staining*

Nucleus pulposus tissues isolated from patients and rats IVDs or nucleus pulposus cells stimulated as described were lysed using a RIPA buffer (pH 7.4) containing a protease inhibitor cocktail (Roche). 10 mg of total protein was used for the Western blots. MMP-1, TIMP-1, aggrecan, collagen II and  $\beta$ -actin were detected by immunoblot with their specific antibodies, respectively. The protein bands were

*P. acnes* induces disc degeneration via MMP-1 and TIMP-1



**Figure 1.** *P. acnes* infection dampened aggrecan and collagen II expression by increasing MMP-1 and decreasing TIMP-1 expression in patients. A, B. Western blot analysis of aggrecan, collagen II, MMP-1 and TIMP-1 expression in nucleus pulposus tissues of patients with or without *P. acnes* IVDs infection. Samples from 10 patients with *P. acnes* IVDs infection were pooled together. C. Quantification of MMP-1 and TIMP-1 mRNA expression in nucleus pulposus tissues of 10 patients with or without *P. acnes* IVDs infection. \* $P < 0.05$ ,  $P$  values were analyzed using Student's  $t$  test. Data were presented as mean  $\pm$  SD from three independent experiments.

quantitatively analyzed using an image analysis system (Quantity One software, Bio-Rad, Hercules, CA, USA).

For MMP-1 and TIMP-1 immunofluorescent staining, nucleus pulposus cells were cultured on glass slides as described and then fixed for 30 min in 4% para-formaldehyde. The NPCs were incubated for 16 h at 4°C with MMP-1 and TIMP-1 antibodies respectively. All images were observed using a fluorescence microscope (Axio, Carl Zeiss, Oberkochen, Germany).

**Real-time quantitative RT-PCR**

Real-time RT-PCR specific primers were used to evaluate gene expression. Rat GAPDH: 5'-TC-TACCCACGGCAAGTCC-3' (forward) and 5'-GATGTTAGCGGGATCTCG-3' (reverse); rat MMP-1: 5'-TTTGATGGACCTCAATAT-3' (forward) and 5'-CATTAGTGCTCCTACA-3' (reverse); rat TIMP-1: 5'-TTTGATCTCTGGCCTCTG-3' (forward) and 5'-CATCTTGATCTCATAACGC-3' (reverse); human GAPDH 5'-CTTAGCACCCCTGGCCAAG-3' (forward) and 5'-TGGTCATGAGTCCCTCCACG-3' (reverse); human MMP-1: 5'-GATGTGGAGTGCCTGA-

TGTG-3' (forward) and 5'-TCTCAATGGCATGGT-CCAC-3' (reverse); rat TIMP-1: 5'-ACATCCGGT-TCGTCTACACC-3' and 5'-TGATGTGCAAGAGTC-CATCC-3' (reverse); RNA analysis was done as previously reported [13].

**Statistical analysis**


All data are representative of three independent experiments and are as mean  $\pm$  S.D. We used two-tailed  $t$  tests to determine significances between the two groups. We did analyses of multiple groups by one-way ANOVA with a Bonferroni post test of GraphPad prism version 5. For all statistical tests, we considered  $P$  value  $< 0.05$  to be statistically significant.

**Results**

***P. acnes* induces IVDD by increasing MMP-1 and decreasing TIMP-1 expression in patients**

To detect the pathological mechanisms of *P. acnes*-induced IVDD, ten *P. acnes*-positive degenerated IVDs were collected from patients; meanwhile, ten *P. acnes*-negative degenerated

*P. acnes* induces disc degeneration via MMP-1 and TIMP-1



**Figure 2.** *P. acnes* infection dampened aggrecan and collagen II expression by increasing MMP-1 and decreasing TIMP-1 expression in rats. A, B. Western blot analysis of aggrecan, collagen II, MMP-1 and TIMP-1 expression in nucleus pulposus tissues of rats with or without *P. acnes* caudal IVDs infection. Samples from 6 rats with *P. acnes* caudal IVDs infection were pooled together. C. Quantification analysis of MMP-1 and TIMP-1 expression in nucleus

## *P. acnes* induces disc degeneration via MMP-1 and TIMP-1

pulposus tissues of 6 rats with or without *P. acnes* caudal IVDs infection. D. Immunofluorescent staining analysis of MMP-1 and TIMP-1 expression in nucleus pulposus tissues of rats with or without *P. acnes* caudal IVDs infection. Scale bar represents 100  $\mu\text{m}$ . \* $P < 0.05$ ,  $P$  values were analyzed using Student's  $t$  test. Data were presented as mean  $\pm$  SD from three independent experiments.

IVDs were chosen as controls with a case-controlled matched method used previously [9, 14]. Consistent with our previous report [11], aggrecan and collagen II expressions were significantly decreased in nucleus pulposus tissues from *P. acnes*-positive degenerated IVDs compared to those from *P. acnes*-negative degenerated IVDs (**Figure 1A, 1B**); from this we predicted that *P. acnes* infection worsened IVDD. Since the concerted activity of matrix-degrading metalloproteinases (MMPs) and aggrecan-degrading enzymes causes a progressive loss of collagens and proteoglycans, and MMP-1 is one of the most important MMPs in ECM degradation [15-18], we then examined whether *P. acnes* dampened aggrecan and collagen II expression by inducing MMP-1 expression. The results showed that *P. acnes* infection strongly increased MMP-1 but decreased the tissue inhibitor of metalloproteinase-1 (TIMP1) expression (**Figure 1A-C**). Taken together, these data demonstrated that *P. acnes* IVDs infection dampened aggrecan and collagen II expression by increasing MMP-1 and decreasing TIMP-1 expression in patients.

### *P. acnes* induces IVDD by increasing MMP-1 and decreasing TIMP-1 expression in rats


To further confirm the dampened aggrecan and collagen II expression resulting from the upregulation of MMP-1 and downregulation of TIMP-1 in *P. acnes*-induced IVDD, *P. acnes* was inoculated into the caudal IVD of rats. We found that *P. acnes* infection significantly decreased aggrecan and collagen II expression in rat nucleus pulposus tissues (**Figure 2A, 2B**). Interestingly, a Western blot (**Figure 2A, 2B**), mRNA quantification (**Figure 2C**), and immunofluorescence analyses (**Figure 2D**) further confirmed the increase in MMP-1 and decrease in TIMP-1 expression in nucleus pulposus tissues in response to *P. acnes* IVDs infection. Collectively, these results demonstrated that *P. acnes* IVDs infection dampened aggrecan and collagen II expression by increasing MMP-1 and decreasing TIMP-1 expression in rats.

### *P. acnes* incubation dampens aggrecan and collagen II expression by increasing MMP-1 and decreasing TIMP-1 expression in NPCs

Next, we investigated whether *P. acnes* infection could dampen aggrecan and collagen II expression by increasing MMP-1 and decreasing TIMP-1 expression *in vitro*. Upon co-culturing NPCs with *P. acnes* (MOI=100) for different time periods (12, 24 h), a time-dependent decrease in aggrecan and collagen II was observed (**Figure 3A, 3B**). Simultaneously, we detected a time-dependent increase in MMP-1 and a decrease in TIMP-1 expression using western blot and real time PCR (**Figure 3A-D**). Consistently, the increased MMP-1 and decreased TIMP-1 expression was further confirmed by immunofluorescence (**Figure 3E**). Taken together, these data revealed that *P. acnes* incubation dampened aggrecan and collagen II expression by increasing MMP-1 and decreasing TIMP-1 expression in NPCs.

### *P. acnes* infection increases MMP-1 and decreased TIMP-1 expression via NF- $\kappa$ B pathway

Having observed that *P. acnes* infection dampened aggrecan and collagen II expression by increasing MMP-1 and decreasing TIMP-1 expression, we next sought to detect the mechanisms by which *P. acnes* infection increases MMP-1 and decreases TIMP-1 expression. Since our previous study demonstrated that the NF- $\kappa$ B signaling pathway mediated *P. acnes*-induced IVDD by regulating aggrecan and collagen II expression, we speculated that NF- $\kappa$ B activation was essential for *P. acnes*-mediated MMP-1 and TIMP-1 expression. To this end, the NF- $\kappa$ B activation was analyzed in response to *P. acnes* infection. The results showed that *P. acnes* incubation increased the phosphorylation of p65 (a subunit of NF- $\kappa$ B) in a time-dependent manner (**Figure 4A, 4B**). Next, we inhibited NF- $\kappa$ B activation with the NF- $\kappa$ B inhibitor-BAY11-7082 and revealed that in line with the dampened p65 phosphorylation, BAY11 pretreatment significantly decreased MMP-1, but it increased TIMP-1 expression in NPCs (**Figure 4C, 4D**). Collectively,



**Figure 3.** *P. acnes* incubation dampened aggrecan and collagen II expression by increasing MMP-1 and decreasing TIMP-1 expression in nucleus pulposus cells. (A, B) Western blot analysis of aggrecan, collagen II, MMP-1 and TIMP-1 expression in nucleus pulposus cells inoculated with or without *P. acnes* (MOI=100) for different times. (C, D) Quantification analysis of MMP-1 (C) and TIMP-1 (D) mRNA expression in nucleus pulposus cells inoculated with or without *P. acnes* (MOI=100) for different times. (E) Immunofluorescent staining analysis of MMP-1 and TIMP-1 expression in nucleus pulposus cells inoculated with or without *P. acnes* (MOI=100) for 24 hours. Scale bar represents 100  $\mu$ m. \* $P < 0.05$ ,  $P$  values were analyzed using one-way ANOVA. Data were presented as mean  $\pm$  SD from three independent experiments.


these data demonstrated that *P. acnes* infection increased MMP-1 and decreased TIMP-1 expression via the NF- $\kappa$ B pathway.

### Discussion

Since the first isolation of low virulence anaerobic bacteria from non-pyogenic degenerated intervertebral discs [5], the pathological roles

of these bacteria in non-pyogenic intervertebral discs have drawn an increasing amount of attention. Accumulating evidence suggests that *P. acnes* is a novel pathogenic factor contributing to intervertebral disc degeneration (IVDD), but the mechanisms by which *P. acnes* induces IVDD are unclear. Here, we found that *P. acnes* infection dampened aggrecan and collagen II synthesis in NPCs by inducing an

*P. acnes* induces disc degeneration via MMP-1 and TIMP-1



**Figure 4.** *P. acnes* infection increased MMP-1 and decreased TIMP-1 expression via NF-κB pathway. A, B. *P. acnes* inoculation (MOI=100) induces p65 phosphorylation in a time-dependent manner. C, D. Western blot analysis of P-p65, MMP-1 and TIMP-1 expression in nucleus pulposus cells induced by *P. acnes* (MOI=100) with or without BAY11-7082 pretreatment. \* $P < 0.05$ ,  $P$  values were analyzed using one-way ANOVA. Data were presented as mean  $\pm$  SD from three independent experiments.

increase in MMP-1 and a decrease in TIMP-1 expression via the NF-κB pathway, which might lead to IVDD.

The most significant biochemical change in IVDD is the loss of the ECM, which mainly contains proteoglycan and collagen II [19, 20]. The matrix metalloproteinases (MMPs) and their inhibitors, the tissue inhibitor of metalloproteinases (TIMPs), are important regulators of ECM turnover [21, 22]. Although our previous study demonstrated that *P. acnes* infection decreased aggrecan and collagen II expression [11], whether MMPs and TIMPs were involved in this system was unknown. In this study, we found that *P. acnes* IVDs infection increased MMP-1 expression, while decreasing TIMP-1 expression, in NPCs both *in vivo* and *in vitro*. Furthermore, our results revealed that NF-κB activation was critical for *P. acnes*-mediated MMP-1 and TIMP-1 expression changes. Although MMP-1 is reported as one of the most

important MMPs in ECM degradation [15-18], MMP-3, MMP-13, ADAMTS4, and ADAM5 have also been implicated in this process; therefore, the possible regulation of the expression of these MMPs and ADAMTSs by *P. acnes* in disc degeneration is a path for further investigation.

The NF-κB pathway is an important catabolic pathway in the pathophysiology of IVDD [23, 24]. In addition, NF-κB has been reported to play a crucial role in the production of catabolic gene expression, such as MMP-1, MMP-13, ADAMTS4, and ADAMTS5 in chondrocytes [25, 26]. However, whether NF-κB was involved in *P. acnes*-mediated MMP-1 expression was unknown. Here, we found that *P. acnes* increased MMP-1 expression and decreased TIMP-1 expression via the NF-κB pathway, to mediate aggrecan and collagen II expression. In addition, our previous study has demonstrated an NF-κB-dependent inflammatory response

in NPCs in response to *P. acnes* infection [11]. Since inflammatory cytokines, such as IL-1 $\beta$ , TNF- $\alpha$ , and IL-6, are involved in mediating MMP-1 and TIMP-1 expression during disc degeneration [27], we speculate that *P. acnes* regulates MMP-1 and TIMP-1 expression through mediating the expression of inflammatory cytokines, but this still needs further examination.

In conclusion, our study provides the first evidence that *P. acnes* IVDDs infection dampens aggrecan and collagen II expression by increasing MMP-1 and decreasing TIMP-1 expression via the NF- $\kappa$ B pathway, which may ultimately lead to IVDD. The confirmation of *P. acnes* as a pathogenic factor for IVDD and elucidation of the underlying mechanisms provides new insights into IVDD and may ultimately lead to the development of novel treatment regimens for IVDD.

#### Acknowledgements

This work was supported by grants from the Shanghai Municipal Health and Family Planning Commission (No. 20164Y0164 and No. 20164Y0132); grants from Ruijin Hospital North, Shanghai Jiaotong University School of Medicine (No. 2015FY03); the Science and Technology Commission of Shanghai Municipality (No. 13430722100 and No. 15DZ19-42604) and from the Shanghai Bureau of Health, Shanghai, China (No. XBR2011024).

#### Disclosure of conflict of interest

None.

**Address correspondence to:** Dr. Changwei Li, Shanghai Key Laboratory for Prevention and Treatment of Bone and Joint Diseases with Integrated Chinese-Western Medicine, Shanghai Institute of Traumatology and Orthopedics, Shanghai Jiaotong University School of Medicine, 197 Ruijin Er Road, Shanghai 200025, China. Tel: +86 21 64313534; Fax: +86 21 64335742; E-mail: changwei393331@163.com; Drs. Xiangyang Xu and Peng Cao, Department of Orthopedics, Ruijin Hospital, Shanghai Jiaotong University School of Medicine, 197 Ruijin Er Road, Shanghai 200025, China. Tel: +86 21 64313534; Fax: +86 21 64335742; E-mail: xu664531@hotmail.com (XYX); dr\_caopeng8@163.com (PC)

#### References

- [1] Le Maitre CL, Pockert A, Buttle DJ, Freemont AJ and Hoyland JA. Matrix synthesis and degradation in human intervertebral disc degeneration. *Biochem Soc Trans* 2007; 35: 652-655.
- [2] Liu Z, Li C, Meng X, Bai Y, Qi J, Wang J, Zhou Q, Zhang W and Zhang X. Hypoxia-inducible factor-1 $\alpha$  mediates aggrecan and collagen Pi expression via NOTCH1 signaling in nucleus pulposus cells during intervertebral disc degeneration. *Biochem Biophys Res Commun* 2017; 488: 554-561.
- [3] Modic MT and Ross JS. Lumbar degenerative disk disease. *Radiology* 2007; 245: 43-61.
- [4] Capoor MN, Ruzicka F, Schmitz JE, James GA, Machackova T, Jancalek R, Smrcka M, Lipina R, Ahmed FS, Alamin TF, Anand N, Baird JC, Bhatia N, Demir-Deviren S, Eastlack RK, Fisher S, Garfin SR, Gogia JS, Gokaslan ZL, Kuo CC, Lee YP, Mavrommatis K, Michu N, Noskova H, Raz A, Sana J, Shamie AN, Stewart PS, Stonemetz JL, Wang JC, Witham TF, Coscia MF, Birkenmaier C, Fischetti VA and Slaby O. *Propionibacterium acnes* biofilm is present in intervertebral discs of patients undergoing microdiscectomy. *PLoS One* 2017; 12: e0174518.
- [5] Stirling A, Worthington T, Rafiq M, Lambert PA and Elliott TS. Association between sciatica and *Propionibacterium acnes*. *Lancet* 2001; 357: 2024-2025.
- [6] Coscia MF, Denys GA and Wack MF. *Propionibacterium acnes*, coagulase-negative staphylococcus, and the "biofilm-like" intervertebral disc. *Spine (Phila Pa 1976)* 2016; 41: 1860-1865.
- [7] Rao PJ, Phan K, Reddy R, Scherman DB, Taylor P and Mobbs RJ. DISC (degenerate-disc infection study with contaminant control): pilot study of Australian cohort of patients without the contaminant control. *Spine (Phila Pa 1976)* 2016; 41: 935-939.
- [8] Albert HB, Lambert P, Rollason J, Sorensen JS, Worthington T, Pedersen MB, Norgaard HS, Vernallis A, Busch F, Manniche C and Elliott T. Does nuclear tissue infected with bacteria following disc herniations lead to Modic changes in the adjacent vertebrae? *Eur Spine J* 2013; 22: 690-696.
- [9] Yuan Y, Zhou Z, Jiao Y, Li C, Zheng Y, Lin Y, Xiao J, Chen Z and Cao P. Histological identification of *propionibacterium acnes* in nonpyogenic degenerated intervertebral discs. *Biomed Res Int* 2017; 2017: 6192935.
- [10] Chen Z, Zheng Y, Yuan Y, Jiao Y, Xiao J, Zhou Z and Cao P. Modic changes and disc degeneration caused by inoculation of *propionibacterium acnes* inside intervertebral discs of rab-

## *P. acnes* induces disc degeneration via MMP-1 and TIMP-1

- bits: a pilot study. *Biomed Res Int* 2016; 2016: 9612437.
- [11] Lin Y, Jiao Y, Yuan Y, Zhou Z, Zheng Y, Xiao J, Li C, Chen Z and Cao P. *Propionibacterium acnes* induces intervertebral disc degeneration by promoting nucleus pulposus cell apoptosis via the TLR2/JNK/mitochondrial-mediated pathway. *Emerg Microbes Infect* 2018; 7: 1.
- [12] O'Donnell MJ, Chin SL, Rangarajan S, Xavier D, Liu L, Zhang H, Rao-Melacini P, Zhang X, Pais P, Agapay S, Lopez-Jaramillo P, Damasceno A, Langhorne P, McQueen MJ, Rosengren A, Dehghan M, Hankey GJ, Dans AL, Elsayed A, Avezum A, Mondo C, Diener HC, Ryglewicz D, Czlonkowska A, Pogosova N, Weimar C, Iqbal R, Diaz R, Yusoff K, Yusufali A, Oguz A, Wang X, Penaherrera E, Lanan F, Ogah OS, Ogunniyi A, Iversen HK, Malaga G, Rumboldt Z, Oveisgharan S, Al Hussain F, Magazi D, Nilanont Y, Ferguson J, Pare G, Yusuf S; INTERSTROKE investigators. Global and regional effects of potentially modifiable risk factors associated with acute stroke in 32 countries (INTERSTROKE): a case-control study. *Lancet* 2016; 388: 761-775.
- [13] Li C, Chen K, Kang H, Yan Y, Liu K, Guo C, Qi J, Yang K, Wang F, Guo L, He C and Deng L. Double-stranded RNA released from damaged articular chondrocytes promotes cartilage degeneration via toll-like receptor 3-interleukin-33 pathway. *Cell Death Dis* 2017; 8: e3165.
- [14] Zhou Z, Chen Z, Zheng Y, Cao P, Liang Y, Zhang X, Wu W, Xiao J and Qiu S. Relationship between annular tear and presence of *Propionibacterium acnes* in lumbar intervertebral disc. *Eur Spine J* 2015; 24: 2496-2502.
- [15] Burrage PS, Mix KS and Brinckerhoff CE. Matrix metalloproteinases: role in arthritis. *Front Biosci* 2006; 11: 529-543.
- [16] Lee YA, Choi HM, Lee SH, Hong SJ, Yang HI, Yoo MC and Kim KS. Hypoxia differentially affects IL-1beta-stimulated MMP-1 and MMP-13 expression of fibroblast-like synoviocytes in an HIF-1alpha-dependent manner. *Rheumatology (Oxford)* 2012; 51: 443-450.
- [17] Vincenti MP and Brinckerhoff CE. Transcriptional regulation of collagenase (MMP-1, MMP-13) genes in arthritis: integration of complex signaling pathways for the recruitment of gene-specific transcription factors. *Arthritis Res* 2002; 4: 157-164.
- [18] Muller-Ladner U and Gay S. MMPs and rheumatoid synovial fibroblasts: Siamese twins in joint destruction? *Ann Rheum Dis* 2002; 61: 957-959.
- [19] Pattappa G, Li Z, Peroglio M, Wismer N, Alini M and Grad S. Diversity of intervertebral disc cells: phenotype and function. *J Anat* 2012; 221: 480-496.
- [20] Adams MA, Dolan P and McNally DS. The internal mechanical functioning of intervertebral discs and articular cartilage, and its relevance to matrix biology. *Matrix Biol* 2009; 28: 384-389.
- [21] Birkedal-Hansen H, Moore WG, Bodden MK, Windsor LJ, Birkedal-Hansen B, DeCarlo A and Engler JA. Matrix metalloproteinases: a review. *Crit Rev Oral Biol Med* 1993; 4: 197-250.
- [22] Gomez DE, Alonso DF, Yoshiji H and Thorgeirsson UP. Tissue inhibitors of metalloproteinases: structure, regulation and biological functions. *Eur J Cell Biol* 1997; 74: 111-122.
- [23] Zhongyi S, Sai Z, Chao L and Jiwei T. Effects of nuclear factor kappa B signaling pathway in human intervertebral disc degeneration. *Spine (Phila Pa 1976)* 2015; 40: 224-232.
- [24] Wang S, Liu C, Sun Z, Yan P, Liang H, Huang K, Li C and Tian J. IL-1beta increases asporin expression via the NF-kappaB p65 pathway in nucleus pulposus cells during intervertebral disc degeneration. *Sci Rep* 2017; 7: 4112.
- [25] Liacini A, Sylvester J, Li WQ, Huang W, Dehnade F, Ahmad M and Zafarullah M. Induction of matrix metalloproteinase-13 gene expression by TNF-alpha is mediated by MAP kinases, AP-1, and NF-kappaB transcription factors in articular chondrocytes. *Exp Cell Res* 2003; 288: 208-217.
- [26] Sandell LJ, Xing X, Franz C, Davies S, Chang LW and Patra D. Exuberant expression of chemokine genes by adult human articular chondrocytes in response to IL-1beta. *Osteoarthritis Cartilage* 2008; 16: 1560-1571.
- [27] Risbud MV and Shapiro IM. Role of cytokines in intervertebral disc degeneration: pain and disc content. *Nat Rev Rheumatol* 2014; 10: 44-56.